arXiv:1606.08349v1 [math.FA] 27 Jun 2016

# A NOTE ON APPROXIMATELY BIFLAT BANACH ALGEBRAS

A. SAHAMI

Abstract. In this paper, we study the notion of approximately biflat Banach algebras for second dual Banach algebras and semigroup algebras. We show that for a locally compact group $G$, if $S(G)^{**}$ is approximately biflat, then $G$ is amenable group. Also we give some conditions which the second dual of a Triangular Banach algebra is never approximately biflat. For a uniformly locally finite semigroup $S$, we show that $\ell^1(S)$ is approximately biflat if and only if $\ell^1(S)$ is biflat.

## 1. Introduction

Helemskii defined the notion of biflat Banach algebras. In fact a Banach algebra $A$ is biflat if there exists a bounded $A$-bimodule $\rho : (A \otimes_p A)^* \to A^*$ such that $\rho \circ \pi_A^*(f) = f$ for each $f \in A^*$, where $\pi_A : A \otimes_p A \to A$ is defined by $\pi_A(a \otimes b) = ab$ for each $a, b \in A$. For a group algebra $L^1(G)$ associated with a locally compact group $G$, $L^1(G)$ is biflat if and only if $G$ is amenable group. For the further details see [6]. Recently Ramsden in [11] characterized the biflatness os semigroup algebras associated to a locally finite inverse semigroup. He showed that for a locally finite inverse semigroup $S$, $\ell^1(S)$ is biflat if and only if each $G_p$ is amenable group, where $p \in E(S)$ and $G_p$ is a maximal subgroup of $S$. Also biflatness of Triangular Banach algebras have been studied in [10].

Approximate notions in the homology of Banach algebras introduced and have been under more observations, recently. A Banach algebra $A$ is approximately biflat if there exists a a net of $A$-bimodule morphisms $(\rho_\alpha)$ from $(A \otimes_p A)^*$ into $A^*$ such that $\rho_\alpha \circ \pi_A^* \xrightarrow{W^*OT} id_{A^*}$, where $W^*OT$ is denoted for the weak-star operator topology and $id_{A^*}$ the identity map on $A^*$. In fact for the discrete Heisenberg group $G$ the Fourier algebra $A(G)$ is approximately biflat but $A(G)$ is not biflat, see [13]. Samei *et al.* also showed that if $A$ is an approximately biflat Banach algebra with an approximate identity, then $A$ is pseudo-amenable, that is, there exists a net $(m_\alpha)$ in $A \otimes_p A$ such that

$$a \cdot m_\alpha - m_\alpha \cdot a, \quad \pi_A(m_\alpha)a \to a \quad (a \in A).$$

For further details about pseudo-amenability, see [5].

Motivated by these considerations, we study approximate biflatness of $\ell^1(S)$, where $S$ is a uniformly locally finite semigroup. We show that $\ell^1(S)$ is approximately biflat if and only

2010 *Mathematics Subject Classification.* Primary 43A07, 20M18, Secondary 46H05.

*Key words and phrases.* Approximate biflatness, Triangular Banach algebras,Segal algebras, semigroup algebras.

if $\ell^1(S)$ is biflat. Also we show that approximately biflatness of $\ell^1(S)^{**}$ implies the pseudo-amenability of $\ell^1(S)$. Also for a locally compact group $G$, we show that approximately biflatness of $S(G)^{**}$, implies that $G$ is amenable, where $S(G)$ is a Segal algebra with respect to $G$. Finally we give a criteria to study approximately biflatness of Triangular Banach algebras. We show that some second dual of Triangular Banach algebras related to a locally compact groups are never approximately biflat.

## 2. Preliminaries

Let $A$ be a Banach algebra. We recall that if $X$ is a Banach $A$-bimodule, then $X^*$ is also a Banach $A$-bimodule via the following actions

$$(a \cdot f)(x) = f(x \cdot a), \quad (f \cdot a)(x) = f(a \cdot x) \quad (a \in A, x \in X, f \in X^*).$$

Throughout, the character space of $A$ is denoted by $\Delta(A)$, that is, all non-zero multiplicative linear functionals on $A$. Let $\phi \in \Delta(A)$. Then $\phi$ has a unique extension $\tilde{\phi} \in \Delta(A^{**})$ which is defined by $\tilde{\phi}(F) = F(\phi)$ for every $F \in A^{**}$.

Let $\{A_\alpha\}_{\alpha\in\Gamma}$ be a collection of Banach algebras. Then we define the $\ell^1$-direct sum of $A_\alpha$ by

$$\ell^1 - \oplus_{\alpha\in\Gamma} A_\alpha = \{(a_\alpha) \in \prod_{\alpha\in\Gamma} A_\alpha : \sum_{\alpha\in\Gamma} ||a_\alpha|| < \infty\}.$$

It is easy to verify that

$$\Delta(\ell^1 - \oplus_{\alpha\in\Gamma} A_\alpha) = \{\oplus\phi_\beta : \phi_\beta \in \Delta(A_\beta), \beta \in \Gamma\},$$

where $\oplus\phi_\beta((a_\alpha)_{\alpha\in\Gamma}) = \phi_\beta(a_\beta)$ for every $(a_\alpha)_{\alpha\in\Gamma} \in \ell^1 - \oplus_{\alpha\in\Gamma} A_\alpha$ and every $\beta \in \Gamma$.

Let $A$ be a Banach algebra and let $\Lambda$ be a non-empty set. The set of all $\Lambda \times \Lambda$ matrixes $(a_{i,j})_{i,j}$ which entries come from $A$ is denoted by $\mathbb{M}_\Lambda(A)$. With the matrix multiplication and the following norm

$$||(a_{i,j})_{i,j}|| = \sum_{i,j} ||a_{i,j}|| < \infty,$$

$\mathbb{M}_\Lambda(A)$ is a Banach algebra. $\mathbb{M}_\Lambda(A)$ belongs to the class of $\ell^1$-Munn algebras. The map $\theta : \mathbb{M}_\Lambda(A) \to A \otimes_p \mathbb{M}_\Lambda(\mathbb{C})$ defined by $\theta((a_{i,j})) = \sum_{i,j} a_{i,j} \otimes E_{i,j}$ is an isometric algebra isomorphism, where $(E_{i,j})$ denotes the matrix unit of $\mathbb{M}_\Lambda(\mathbb{C})$. Also it is well-known that $\mathbb{M}_\Lambda(\mathbb{C})$ is a biprojective Banach algebra [11, Proposition 2.7].

The main reference for the semigroup theory is [7]. Let $S$ be a semigroup and let $E(S)$ be the set of its idempotents. A partial order on $E(S)$ is defined by

$$s \leq t \Longleftrightarrow s = st = ts \quad (s, t \in E(S)).$$

If $S$ is an inverse semigroup, then there exists a partial order on $S$ which is coincide with the partial order on $E(S)$. Indeed

$$s \leq t \Longleftrightarrow s = ss^*t \quad (s, t \in S).$$

For every $x \in S$, we denote $(x] = \{y \in S | \, y \leq x\}$. $S$ is called locally finite (uniformly locally finite) if for each $x \in S$, $|(x]| < \infty$ ($\sup\{|(x]| : x \in S\} < \infty$), respectively.

Suppose that $S$ is an inverse semigroup. Then the maximal subgroup of $S$ at $p \in E(S)$ is denoted by $G_p = \{s \in S | ss^* = s^*s = p\}$. For an inverse semigroup $S$ there exists a relation $\mathfrak{D}$ such that $s\mathfrak{D}t$ if and only if there exists $x \in S$ such that $ss^* = xx^*$ and $t^*t = x^*x$. We denote $\{\mathfrak{D}_\lambda : \lambda \in \Lambda\}$ for the collection of $\mathfrak{D}$-classes and $E(\mathfrak{D}_\lambda) = E(S) \cap \mathfrak{D}_\lambda$. An inverse semigroup $S$ is called Clifford if for each $s \in S$, there exists $s^* \in S$ such that $ss^* = s^*s$.

## 3. Approximate biflatness of second dual of Banach algebras

In this section we investigate approximate biflatness dual Banach algebras.

**Proposition 3.1.** *Let $A$ be a Banach algebra. $A$ is approximately biflat Banach algebra if and only if there exists a net $(\rho_\alpha)$ of bounded $A$-bimodule morphisms from $A$ into $(A \otimes_p A)^{**}$ such that $\pi_A^{**} \circ \rho_\alpha(a) \to a$ for every $a \in A$.*

*Proof.* Let $A$ be approximately biflat. Then there exists a net $\xi_\alpha : A^* \to (A \otimes_p A)^*$ such that $\xi_\alpha \circ \pi_A^*(f) - f \to 0$, for every $f \in A^*$. Set $\rho_\alpha = \xi_\alpha^*$, hence for each $a \in A$ and $f \in A^*$ with $||f|| \leq 1$, we have

$$\begin{aligned} ||\pi_A^{**} \circ \rho_\alpha(a) - a|| = ||\pi_A^{**} \circ \xi_\alpha^*(a) - a|| &= ||(\xi_\alpha \circ \pi_A^*)^*(a) - a|| \\ &= ||(\xi_\alpha \circ \pi_A^*)^*(a)(f) - a(f)|| \\ &= ||a(\xi_\alpha \circ \pi_A^*(f)) - a(f)|| \\ &= ||a(\xi_\alpha \circ \pi_A^*(f) - f)|| \\ &\leq ||a||||\xi_\alpha \circ \pi_A^*(f) - f|| \to 0. \end{aligned}$$

For converse, suppose that there exists a net $(\rho_\alpha)$ of bounded $A$-bimodule morphisms from $A$ into $(A \otimes_p A)^{**}$ such that $\pi_A^{**} \circ \rho_\alpha(a) \to a$ for every $a \in A$. Set $\xi_\alpha = \rho_\alpha^*|_{(A \otimes_p A)^*}$.

$$\xi_\alpha \circ \pi_A^*(f)(a) - f(a) = f(\pi_A^{**} \circ \rho_\alpha(a) - a) \to 0,$$

where $f \in A^*, a \in A$. □

**Theorem 3.2.** *Let $A$ be a Banach algebra with an approximate identity. If $A^{**}$ is approximately biflat, then $A$ is pseudo-amenable.*

*Proof.* Suppose that $A^{**}$ is approximately biflat. Then by Proposition 3.1 there exists a net $(\rho_\alpha)_{\alpha \in I}$ of $A^{**}$-bimodule morphism from $A^{**}$ into $(A^{**} \otimes_p A^{**})^{**}$ such that $\pi_{A^{**}}^{**} \circ \rho_\alpha(a) \to a$ for each $a \in A^{**}$. It is easy to see that the net $(\rho_\alpha|_A)$ is also a net of $A$-bimodule morphism satisfies $\pi_{A^{**}}^{**} \circ \rho_\alpha|_A(a) \to a$ for each $a \in A$. We denote $(e_\lambda)_{\lambda \in J}$ for the approximate identity of $A$. Consider

$$\lim_\alpha \lim_\lambda a \cdot \rho_\alpha(e_\lambda) - \rho_\alpha(e_\lambda) \cdot a = \lim_\alpha \lim_\lambda \rho_\alpha(ae_\lambda - e_\lambda a) = \lim_\alpha \rho_\alpha(0) = 0 \quad (a \in A).$$

Also

$$\lim_{\alpha}\lim_{\lambda}\pi^{**}_{A^{**}}\circ\rho_{\alpha}(e_{\lambda})a-a=\lim_{\alpha}\pi^{**}_{A^{**}}\circ\rho_{\alpha}(a)-a=0,\quad (a\in A).$$

Let $E=I\times J^I$ be a directed set with product ordering, that is

$$(\alpha,\beta)\leq_E(\alpha^{'},\beta^{'})\Leftrightarrow\alpha\leq_I\alpha^{'},\beta\leq_{J^I}\beta^{'}\qquad(\alpha,\alpha^{'}\in I,\quad\beta,\beta^{'}\in J^I),$$

where $J^I$ is the set of all functions from $I$ into $J$ and $\beta\leq_{J^I}\beta^{'}$ means that $\beta(d)\leq_J\beta^{'}(d)$ for each $d\in I$. Suppose that $\gamma=(\alpha,\beta_{\alpha})\in E$ and $m_{\gamma}=\rho_{\alpha}(e_{\lambda_{\alpha}})\in(A^{**}\otimes_p A^{**})^{**}$. Applying iterated limit theorem [9, page 69] and above calculations, we can easily see that

$$a\cdot m_{\gamma}-m_{\gamma}\cdot a\to 0,\quad \pi^{**}_{A^{**}}(m_{\gamma})a\to a,\quad (a\in A).$$

There exists a bounded linear map $\psi:A^{**}\otimes_p A^{**}\to(A\otimes_p A)^{**}$ such that for $a,b\in A$ and $m\in A^{**}\otimes_p A^{**}$, the following holds;

(i) $\psi(a\otimes b)=a\otimes b$,
(ii) $\psi(m)\cdot a=\psi(m\cdot a),\qquad a\cdot\psi(m)=\psi(a\cdot m)$,
(iii) $\pi^{**}_A(\psi(m))=\pi_{A^{**}}(m)$,

see [4, Lemma 1.7]. So $\psi^{**}(m_{\gamma})$ is a net in $(A\otimes_p A)^{****}$ such that

$$a\cdot\psi^{**}(m_{\gamma})-\psi^{**}(m_{\gamma})\cdot a\to 0,\quad \pi^{****}_A(\psi^{**}(m_{\gamma}))a=\pi^{**}_{A^{**}}(m_{\gamma})a\to a\qquad(a\in A).$$

Put $n_{\gamma}=\psi^{**}(m_{\gamma})$. Suppose that $\epsilon>0$ and $F=\{a_1,...,a_r\}\subseteq A$. Set

$$\begin{aligned}V=&\{(a_1\cdot n-n\cdot a_1,...,a_r\cdot n-n\cdot a_r,\pi^{**}_A(n)a_1-a_1,...,\pi^{**}_A(n)a_r-a_r)|n\in(A\otimes_p A)^{**}\}\\&\subseteq(\prod_{i=1}^{r}(A\otimes_p A)^{**})\oplus_1(\prod_{i=1}^{r}A^{**}).\end{aligned}$$

It is easy to see that $(0,0,...,0)$ is a $w$-limit point of $V$. Since $V$ is convex set $\overline{V}^{||\cdot||}=\overline{V}^{w}$, then $(0,0,...,0)$ is a $||\cdot||$-limit point of $V$. Hence there exists a net $(n_{(F,\epsilon)})$ in $(A\otimes_p A)^{**}$ such that

$$||a_i\cdot n_{(F,\epsilon)}-n_{(F,\epsilon)}\cdot a_i||<\epsilon,\quad ||\pi^{**}_A(n_{(F,\epsilon)})a_i-a_i||<\epsilon,\quad(i\in\{1,2,...,r\}).$$

Observe that

$$\Delta=\{(F,\epsilon):F\text{ is a finite subset of }A,\epsilon>0\},$$

with the following order

$$(F,\epsilon)\leq(F',\epsilon')\Longrightarrow F\subseteq F',\quad\epsilon\geq\epsilon'$$

is a directed set. It is easy to see that there exists a net $(n_{(F,\epsilon)})_{(F,\epsilon)\in\Delta}$ in $(A\otimes_p A)^{**}$ such that

$$a\cdot n_{(F,\epsilon)}-n_{(F,\epsilon)}\cdot a\to 0,\quad\pi^{**}_A(n_{(F,\epsilon)})a-a\to 0,$$

for every $a\in A$. Using the same method as above we can assume that $(n_{(F,\epsilon)})_{(F,\epsilon)\in\Delta}$ is a subset of $A\otimes_p A$. This means that $A$ is pseudo-amenable. □

Let $A$ be a Banach algebra and $\phi \in \Delta(A)$. We say that $A$ is approximately $\phi$-inner amenable if there exists a net $(a_\alpha)_\alpha$ in $A$ such that $aa_\alpha - a a_\alpha a \to 0$ and $\phi(a_\alpha) \to 1$, for all $a \in A$. Also $A$ is approximately left $\phi-$amenable if there exists a net $m_\alpha$ in $A$ such that $am_\alpha - \phi(a)m_\alpha \to 0$ and $\phi(m_\alpha) \to 1$, see [1].

The proof of following two results are similar to the proof of Theorem 3.2 which we omit it.

**Theorem 3.3.** *Suppose that $A$ is an approximately $\phi$-inner amenable Banach algebra. If $A^{**}$ is approximately biflat, then $A$ is approximately left $\phi$-amenable.*

**Corollary 3.4.** *Suppose that $A$ is an approximately $\phi$-inner amenable Banach algebra. If $A$ is approximately biflat, then $A$ is approximately left $\phi$-amenable.*

**Theorem 3.5.** *Let $A$ be a Banach algebra with $\phi \in \Delta(A)$. Suppose that $\overline{Aker\phi} = ker\phi$. If $A^{**}$ is approximately biflat, then $A$ is left $\phi$-amenable.*

*Proof.* Since $A^{**}$ is approximately biflat, there exists a net of $A^{**}$-bimodule morphisms $(\rho_\alpha)$ from $A^{**}$ into $(A^{**} \otimes_p A^{**})^{**}$ such that $\pi^{**}_{A^{**}} \circ \rho_\alpha(a) \to a \quad (a \in A^{**})$. We denote $id : A \to A$ for the identity map and $q : A \to \frac{A}{L}$ the quotient map, where $L = ker\phi$. Also it is well-known that there exists a bounded linear map $\psi : A^{**} \otimes_p A^{**} \to (A \otimes_p A)^{**}$ such that for $a, b \in A$ and $m \in A^{**} \otimes_p A^{**}$, the following holds;

(i) $\psi(a \otimes b) = a \otimes b$,
(ii) $\psi(m) \cdot a = \psi(m \cdot a), \qquad a \cdot \psi(m) = \psi(a \cdot m)$,
(iii) $\pi^{**}_A(\psi(m)) = \pi_{A^{**}}(m)$,

see [4, Lemma 1.7]. Set $\eta_\alpha := (id \otimes q)^{****} \circ \psi^{**} \circ \rho_\alpha|_A : A \to (A \otimes_p \frac{A}{L})^{****}$ for each $\alpha$. We claim that $\eta_\alpha(l) = 0$ for each $l \in \ker \phi$. To see this let $l \in \ker \phi$ be an arbitrary element. Since $\overline{AL} = L$, there exist two nets $(a_\beta)$ in $A$ and $(l_\beta)$ in $L$ such that $l = \lim_\beta a_\beta l_\beta$. Consider

$$\begin{aligned} \eta_\alpha(l) = (id \otimes q)^{****} \circ \psi^{**} \circ \rho_\alpha(l) &= (id \otimes q)^{****} \circ \psi^{**} \circ \rho_\alpha(l) \\ &= \lim_\beta (id \otimes q)^{****} \circ \psi^{**} \circ \rho_\alpha(a_\beta l_\beta) \\ &= \lim_\beta (id \otimes q)^{****}(\psi^{**} \circ \rho_\alpha(a_\beta) \cdot l_\beta) = 0. \end{aligned} \tag{3.1}$$

Hence, for each $\alpha$, $\eta_\alpha$ can induce a map on $\frac{A}{L}$ which we again denote it by $\eta_\alpha$. We also denote $\overline{\phi}$ for a character which induced by $\phi$ on $\frac{A}{L}$ given by

$$\overline{\phi}(a + L) = \phi(a) \quad (a \in A).$$

Set

$$g_\alpha = (id \otimes \overline{\phi})^{****} \circ \eta_\alpha : \frac{A}{L} \to A^{**}.$$

Pick an element $x_0 \in A$ such that $\phi(x_0) = 1$. Define $m_\alpha = g_\alpha(x_0 + L)$. We know that $(g_\alpha)$ is a net of left $A$-module morphisms. Thus

$$\begin{aligned} am_\alpha = a(id \otimes \overline{\phi})^{****} \circ \eta_\alpha(x_0 + L) &= (id \otimes \overline{\phi})^{****} \circ \eta_\alpha(ax_0 + L) \\ &= \phi(a)(id \otimes \overline{\phi})^{****} \circ \eta_\alpha(x_0 + L) \\ &= \phi(a)m_\alpha, \end{aligned} \tag{3.2}$$

the last equality holds because $ax_0 - \phi(a)x_0 \in L$. Since

$$\widetilde{\overline{\phi}} \circ (id \otimes \overline{\phi})^{****} = (\phi \otimes \overline{\phi})^{****}, \quad (\phi \otimes \overline{\phi})^{****} \circ (id \otimes q)^{****} = \widetilde{\overline{\phi}} \circ \pi_A^{****},$$

we have

$$\begin{aligned} \widetilde{\overline{\phi}}(m_\alpha) = \widetilde{\overline{\phi}} \circ (id \otimes \overline{\phi})^{****} \circ \eta_\alpha(x_0 + L) &= \widetilde{\overline{\phi}} \circ (id \otimes \overline{\phi})^{****} \circ \eta_\alpha(x_0) \\ &= \widetilde{\overline{\phi}} \circ (id \otimes \overline{\phi})^{****} \circ (id \otimes q)^{****} \circ \rho_\alpha(x_0) \\ &= \widetilde{\overline{\phi}} \circ \pi_A^{****} \circ \rho_\alpha(x_0) \\ &\to \phi(x_0) = 1. \end{aligned} \tag{3.3}$$

Replacing $(m_\alpha)$ with $(\frac{m_\alpha}{\phi(m_\alpha)})$ on can find an element $m \in A^{****}$ such that $am = \phi(a)m$ and $\widetilde{\overline{\phi}}(m) = 1$ for every $a \in A$. Let $F = \{a_1, a_2, ..., a_r\}$ be an arbitrary finite subset of $A$ and $\epsilon > 0$. Set

$$V = \{(a_1 n - \phi(a_1)n, a_2 n - \phi(a_2)n, ..., a_r n - \phi(a_r)n, \widetilde{\phi}(n) - 1) | n \in A^{**}, ||n|| \leq ||m||\}.$$

It is easy to see that $V$ is a convex subset of $\prod_{i=1}^r A^{**} \oplus_1 \mathbb{C}$. It is easy to see that $(0, 0, ..., 0) \in \overline{V}^w = \overline{V}^{||\cdot||}$. Thus there exists a bounded net $(n_{(F,\epsilon)})_{(F,\epsilon)}$ in $A^{**}$ such that

$$||a_i n_{(F,\epsilon)} - \phi(a_i)n_{(F,\epsilon)}|| < \epsilon, \quad |\tilde{\phi}(n_{(F,\epsilon)}) - 1| < \epsilon, \quad a_i \in F.$$

It is easy to see that

$$\Delta = \{(F, \epsilon) : F \text{ is a finite subset of } A, \epsilon > 0\},$$

with the following order

$$(F, \epsilon) \leq (F', \epsilon') \Longrightarrow F \subseteq F', \quad \epsilon \geq \epsilon'$$

is a directed set. Therefore there exists a bounded net $(n_{(F,\epsilon)})_{(F,\epsilon)\in\Delta}$ in $A^{**}$ such that

$$an_{(F,\epsilon)} - \phi(a)n_{(F,\epsilon)} \to 0, \quad \tilde{\phi}(n_{(F,\epsilon)}) - 1 \to 0, \quad a \in A.$$

Since $(n_{(F,\epsilon)})_{(F,\epsilon)}$ is a bounded net in $A^{**}$, then $(n_{(F,\epsilon)})_{(F,\epsilon)}$ has a $w^*$-limit point in $A^{**}$, say $N$. It is easy to see that

$$aN = \phi(a)N, \quad \tilde{\phi}(N) = 1 \quad (a \in A).$$

It means that $A$ is left $\phi$-amenable. □

The map $\phi_1 : L^1(G) \to \mathbb{C}$ which specified by

$$\phi_1(f) = \int_G f(x)dx$$

is called augmentation character. We know that augmentation character induce a character on $S(G)$ which we denote it by $\phi_1$ again, see [2].

We recall that, for a locally compact group $G$, a linear subspace $S(G)$ of $L^1(G)$ is said to be a Segal algebra on $G$ if it satisfies the following properties:

(i) $S(G)$ is a dense left ideal in $L^1(G)$;
(ii) $S(G)$ with respect to some norm $||\cdot||_{S(G)}$ is a Banach space and $||f||_{L^1(G)} \leq ||f||_{S(G)}$;
(iii) For $f \in S(G)$ and $y \in G$, $L_y f \in S(G)$ and the map $y \mapsto \delta_y * f$ is continuous. Also $||\delta_y * f||_{S(G)} = ||f||_{S(G)}$, for $f \in S(G)$ and $y \in G$.

For more information about this algebras see [12].

**Corollary 3.6.** *Let $G$ be a locally compact group. If $S(G)^{**}$ is approximately biflat, then $G$ is amenable.*

*Proof.* It is well-known that every Segal algebra has a left approximate identity. Suppose that $\phi \in \Delta(S(G))$. It is easy to see that $\overline{S(G)\ker\phi} = \ker\phi$. Using the Theorem 3.5, approximate biflatness of $S(G)^{**}$ implies that $S(G)$ is left $\phi$-amenable. Now by [2, Corollary 3.4] $G$ is amenable. □

## 4. Approximate biflatness of certain semigroup algebras

In this section we study approximate biflatness of some semigroup algebras.
Before giving the following proposition we have to give some backgrounds. Suppose that $A$ and $B$ are Banach algebras and also suppose that $E$ and $F$ are Banach $A$-bimodule and Banach $B$-bimodule, respectively. Via the following module actions, one can see that $E \otimes_p F$ becomes a Banach $A \otimes_p B$-bimodule:

$$(a \otimes b) \cdot (x \otimes y) = (a \cdot x) \otimes (b \cdot y), \quad (x \otimes y) \cdot (a \otimes b) = (x \cdot a) \otimes (y \cdot b),$$

for each $a \in A, x \in E, b \in B, y \in F$. One can readily see that $B(E, F)$ (the set of all bounded linear operator from $E$ into $F$) is a Banach $A \otimes_p B$-bimodule via the following actions:

$$((a \otimes b) * T)(x) = b \cdot T(x \cdot a), \quad (T * (a \otimes b))(x) = T(a \cdot x) \cdot b,$$

for each $T \in B(E, F), a \in A, b \in B, x \in E$. We denote this Banach $A \otimes_p B$-bimodule by $\tilde{B}(E, F)$. Also we can see that $B(F, E)$ becomes a Banach $A \otimes_p B$-bimodule via the following actions:

$$((a \otimes b) * T)(x) = a \cdot T(x \cdot b), T * (a \otimes b))(x) = T(b \cdot x) \cdot a,$$

for each $T \in B(E, F), a \in A, b \in B, x \in E$. We denote this Banach $A \otimes_p B$-bimodule by $\widehat{B}(E, F)$. Note that for each $\lambda \in (E \otimes_p F)^*$ we can define $\widetilde{T_\lambda} \in B(E, F^*)$ and $\widehat{T_\lambda} \in B(F, E^*)$ by

$$< y, \widetilde{T_\lambda}(x) > = < x \otimes y, \lambda >, \quad < x, \widehat{T_\lambda}(y) > = < x \otimes y, \lambda >,$$

for each $x \in E, y \in F$. The map $\widetilde{\xi} : (E \otimes_p F)^* \to \tilde{B}(E, F^*)$ given by $\widetilde{\xi}(\lambda) = \widetilde{T}_\lambda$ is an isometric $A \otimes_p B$-bimodule isomorphism. Also the map $\widehat{\xi} : (E \otimes_p F)^* \to \widehat{B}(F, E^*)$ given by $\widehat{\xi}(\lambda) = \widehat{T_\lambda}$ is an isometric $A \otimes_p B$-bimodule isomorphism. Then there exists a bounded isometric $A \otimes_p B$-bimodule isomorphism from $\widetilde{B}(E, F^*)$ into $\widehat{B}(F, E^*)$ which denoted by $L$. Also we have to remind that there exists an isometric $A \otimes_p B$-bimodule isomorphism from $(A \otimes_p A) \otimes_p (B \otimes_p B)$ into $(A \otimes_p B) \otimes_p (A \otimes_p B)$ defined by

$$\theta(a \otimes a' \otimes b \otimes b') = a \otimes b \otimes a' \otimes b' \quad (a, a' \in A, b, b' \in B).$$

It is easy to see that $\theta$ is a bounded $A \otimes_p B$-bimodule morphism.

**Proposition 4.1.** *Let $A$ be a biflat Banach algebra and $B$ be approximate biflat Banach algebra. Then $A \otimes_p B$ is approximate biflat.*

*Proof.* Since $A$ is approximate biflat, there exists a net of bounded $A$-bimodules from $A$ into $(A \otimes_p A)^{**}$, say $(\rho_\alpha)_{\alpha \in I}$, such that $\rho_\alpha \circ \pi_A^*(f) - f \to 0$, for each $f \in A^*$. Also since $B$ is biflat, there exists a bounded $B$-bimodules from $B$ into $(B \otimes_p B)^{**}$, say $\rho$, such that $\rho \circ \pi_B^*(g) = g$ for each $g \in B^*$. Set

$$\begin{aligned} \bar{\rho}_\alpha : ((A \otimes_p B) \otimes_p (A \otimes_p B))^* &\xrightarrow{\theta^*} ((A \otimes_p A) \otimes_p (B \otimes_p B))^* \\ &\xrightarrow{\widetilde{\xi}} \widetilde{B}(A \otimes_p A, (B \otimes_p B)^*) \\ &\xrightarrow{T \mapsto \rho \circ T} \widetilde{B}(A \otimes_p A, B^*) \\ &\xrightarrow{L} \widehat{B}(B, (A \otimes_p A)^*) \\ &\xrightarrow{T \mapsto \rho_\alpha \circ T} \widehat{B}(B, A^*) \\ &\xrightarrow{\widehat{\xi^{-1}}} (A \otimes_p B)^*. \end{aligned} \tag{4.1}$$

Since $\bar{\rho}_\alpha$ is a composition of some $(A \otimes_p B)$-bimodule morphisms, then $\bar{\rho}_\alpha$ is a net of $(A \otimes_p B)$-bimodule morphisms. Take $\lambda \in (A \otimes_p B)^*$. Using the following facts

$$\rho \circ \pi_B^* \circ \widetilde{T}_\lambda \circ \pi_A = \widetilde{T}_\lambda \circ \pi_A$$

and

$$L \circ \widetilde{T}_\lambda \circ \pi_A = \pi_A^* \circ \widehat{T}_\lambda$$

we have

$$\overline{\rho}_\alpha \circ \pi^*_{A \otimes_p B}(\lambda) - \lambda = \widehat{\xi^{-1}} \circ \rho_\alpha \circ \pi_A^* \circ \widehat{T_\lambda} - \lambda \to \lambda - \lambda = 0.$$

This finishes the proof. □

We partially shows the converse of above Proposition in the following theorem.

**Theorem 4.2.** *Let $A$ and $B$ be Banach algebras. Suppose that $A$ has an identity and $B$ has a non-zero idempotent. If $A \otimes_p B$ is approximately biflat, then $A$ is approximately biflat, so $A$ is pseudo-amenable.*

*Proof.* Suppose that $A \otimes_p B$ is approximately biflat. Then by Proposition 3.1 there exists a net $(\rho_\alpha)$ of $A$-bimodule morphisms from $A \otimes_p B$ into $((A \otimes_p B) \otimes_p (A \otimes_p B))^{**}$ such that $\pi^{**}_{A\otimes_p B} \circ \rho_\alpha(x) \to x$ for each $x \in A \otimes_p B$. Take $e \in A$ the identity and $b_0 \in B$ the non-zero identity. It is easy to see that $A \otimes_p B$ becomes a Banach $A$-bimodule via the following actions:

$$a_1 \cdot (a_2 \otimes b) = a_1 a_2 \otimes b, \quad (a_2 \otimes b) \cdot a_1 = a_2 a_1 \otimes b \qquad (a_1, a_2 \in A, b \in B).$$

For each $\alpha$, we have

$$\begin{aligned} \rho_\alpha(a_1 a_2 \otimes b_0) = \rho_\alpha((a_1 \otimes b_0)(a_2 \otimes b_0)) &= (a_1 \otimes b_0) \cdot \rho_\alpha(a_2 \otimes b_0) \\ &= (a_1 \cdot (e \otimes b_0)) \cdot \rho_\alpha(a_2 \otimes b_0) \\ &= a_1 \cdot \rho_\alpha(e a_2 \otimes b_0 b_0) \\ &= a_1 \cdot \rho_\alpha(a_2 \otimes b_0). \end{aligned} \tag{4.2}$$

For each $\alpha$, we can also see that

$$\rho_\alpha((a_2 \otimes b_0) \cdot a_1) = \rho_\alpha(a_2 \otimes b_0) \cdot a_1.$$

For each $\alpha$ set $\overline{\rho}_\alpha(a) = \rho_\alpha(a \otimes b_0)$. It is easy to see that $(\overline{\rho}_\alpha)$ is a net of $A$-bimodule morphisms. Since $b_0$ is a non-zero element in $B$, by Hahn-Banach theorem there exists a functional $f \in B^*$ such that $f(b_0) = 1$. Define $T : (A \otimes_p B) \otimes_p (A \otimes_p B) \to A \otimes_p A$ by

$$T(a \otimes b \otimes c \otimes d) = f(bd)a \otimes c$$

for each $a, c \in A$ and $b, d \in B$. Clearly $T$ is a bounded linear map. One can see that $\pi^{**}_A \circ T^{**} = (id_A \otimes f)^{**} \circ \pi^{**}_{A\otimes_p B}$, where

$$id_A \otimes f(a \otimes b) = f(b)a \quad (a \in A, b \in B).$$

Set $\widetilde{\rho}_\alpha = T^{**} \circ \overline{\rho}_\alpha$. It is easy to see that $(\widetilde{\rho}_\alpha)_\alpha$ is a net of bounded $A$-bimodule morphisms. Since $f(b_0) = 1$, we have

$$\begin{aligned} \pi^{**}_A \circ \widetilde{\rho}_\alpha(a) = \pi^{**}_A \circ T^{**} \circ \overline{\rho}_\alpha(a) &= (\pi_A \circ T)^{**} \circ \rho_\alpha(a \otimes b_0) \\ &= (id_A \otimes f)^{**} \circ \pi^{**}_{A\otimes_p B} \cdot \rho_\alpha(a \otimes b_0) \\ &\to (id_A \otimes f)^{**}(a \otimes b_0) = a, \end{aligned} \tag{4.3}$$

for each $a \in A$. Using Proposition 3.1 $A$ is approximate biflat. Since $A$ has an identity by [13, Theorem 2.4], $A$ is pseudo-amenable. □

**Corollary 4.3.** *Let $A$ be a Banach algebra. If $A$ is approximately biflat, then $\mathbb{M}_\Lambda(A)$ is approximately biflat. Converse is true, provided that $A$ has a unit.*

*Proof.* Let $A$ be approximately biflat. It is well-known that there exists an isometrical isomorphism between $\mathbb{M}_\Lambda(A)$ and $A \otimes_p \mathbb{M}_\Lambda(\mathbb{C})$. Using the fact that $\mathbb{M}_\Lambda(\mathbb{C})$ is always biprojective, see [11, Proposition 2.7], then $\mathbb{M}_\Lambda(\mathbb{C})$ is biflat. Now by Proposition 4.1, $A \otimes_p \mathbb{M}_\Lambda(\mathbb{C})$ is approximately biflat.

For converse, since $A$ is unital and $\mathbb{M}_\Lambda(\mathbb{C})$ has a non-zero idempotent, then by Theorem 4.2 the approximate biflatness of $\mathbb{M}_\Lambda(A) \cong A \otimes_p \mathbb{M}_\Lambda(\mathbb{C})$ implies approximate biflatness of $A$. □

**Theorem 4.4.** *Let $S$ be an inverse semigroup such that $E(S)$ is uniformly locally finite. $\ell^1(S)$ is approximately biflat if and only if $\ell^1(S)$ is biflat.*

*Proof.* Suppose that $\ell^1(S)$ is approximately biflat. Then by Proposition 3.1 there exists a bet of $\ell^1(S)$-bimodule morphism from $\ell^1(S)$ into $(\ell^1(S) \otimes_p \ell^1(S))^{**}$ such that $\pi^{**}_{\ell^1(S)} \circ \rho_\alpha(a) - a \to 0$ for each $a \in \ell^1(S)$. Since $S$ is uniformly locally finite, by [11, Theorem 2.18] we have

$$\ell^1(S) \cong \ell^1 - \bigoplus \{\mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda}))\},$$

where is a $\mathfrak{D}$-class and $G_{p_\lambda}$ is a maximal subgroup at $p_\lambda$. Then the map $P_{p_\lambda} : \ell^1(S) \to \mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda}))$ is a continuous homomorphism with a dense range. Define

$$\eta_\alpha : \mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda})) \to (\mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda})) \otimes_p \mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda})))^{**}$$

by $\eta_\alpha = (P \otimes P)^{**} \circ \rho_\alpha|_{\mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda}))}$. It is easy to see that $(\eta_\alpha)$ is a net of $\mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda}))$-bimodule morphisms. For each $a \in \mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda}))$ we have

$$\begin{aligned} \pi^{**}_{\mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda}))} \circ \eta_\alpha(a) &= \pi^{**}_{\mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda}))} \circ (P_{p_\lambda} \otimes P_{p_\lambda})^{**} \circ \rho_\alpha|_{\mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda}))}(a) \\ &= P^{**}_{p_\lambda} \circ \pi^{**}_{\ell^1(S)} \circ \rho_\alpha|_{\mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda}))}(a) \to a. \end{aligned} \tag{4.4}$$

Hence $\mathbb{M}_{E(\mathfrak{D}_\lambda)}(\ell^1(G_{p_\lambda}))$ is approximately biflat. By Theorem 4.2, $\ell^1(G_{p_\lambda})$ is approximately biflat. Since $\ell^1(G_{p_\lambda})$ is unital, by [13, Theorem 2.4] $\ell^1(G_{p_\lambda})$ is pseudo-amenable, hence by [5, Proposition 4.1] $G_{p_\lambda}$ is amenable. Applying [3, Theorem 3.7] to finish the proof.

Converse is clear. □

**Corollary 4.5.** *Let $S = \cup_{e \in E(S)} G_e$ be a Clifford semigroup such that $E(S)$ is uniformly locally finite. If $\ell^1(S)^{**}$ is approximately biflat, then $G_e$ is amenable for each $e \in E(S)$.*

*Proof.* Suppose that $\ell^1(S)^{**}$ is approximately biflat. It is well-known that $\ell^1(S) \cong \ell^1 - \oplus_{e \in E(S)} \ell^1(G_e)$. Since $\ell^1(G_e)$ is unital, then $\ell^1(S)$ has an approximate identity. Applying previous Theorem, implies that $\ell^1(S)$ is pseudo-amenable. Then by [3, Theorem 3.7] $G_e$ is amenable, for each $e \in E(S)$. □

## 5. An application to Triangular Banach algebras

In this section we give some examples of matrix algebras which is never approximately biflat.

Let $A$ be a Banach algebra and $X$ be a Banach $A$-bimodule. Suppose that $\phi \in \Delta(A)$. We say that $X$ has a left $\phi$-character if there exists a non-zero map $\psi \in X^*$ such that

$$\psi(a \cdot x) = \phi(a)\psi(x), \quad (a \in A, x \in X).$$

Similarly we can define right case and two sided case. It is easy to see that for $\phi \in \Delta(A)$, $\phi \otimes \phi$ on $A \otimes_p A$ is a left $\phi$-character. Also if $A$ has a closed ideal $I \subseteq \ker\phi$, then $\overline{\phi}$ on $\frac{A}{I}$ is a left $\phi$-character, where $\overline{\phi} : \frac{A}{L} \to \mathbb{C}$ given by $\overline{\phi}(a + I) = \phi(a)$ for all $a \in A$.

Let $A$ and $B$ be Banach algebras and let $X$ be a Banach $A, B$-module, that is, $X$ is a Banach space, a left $A$-module and a right $B$-module which the compatible module action that satisfies $(a \cdot x) \cdot b = a \cdot (x \cdot b)$ and $||a \cdot x \cdot b|| \leq ||a||||x||||b||$ for every $a \in A, x \in X, b \in B$. With the usual matrix operation and $|| \begin{pmatrix} a & x \\ 0 & b \end{pmatrix} || = ||a|| + ||x|| + ||b||$, $T = Tri(A, X, B) = \begin{pmatrix} A & X \\ 0 & B \end{pmatrix}$ becomes a Banach algebra which is called Triangular Banach algebra. Let $\phi \in \Delta(B)$. We define a character $\psi_\phi \in \Delta(T)$ via $\psi_\phi \begin{pmatrix} a & x \\ 0 & b \end{pmatrix} = \phi(b)$ for every $a \in A$, $b \in B$ and $x \in X$.

**Theorem 5.1.** *Let $T = Tri(A, X, B)$ with $\overline{A^2} = A$ and $\overline{A \cdot X} = X$. Suppose that $\phi \in \Delta(B)$ with $\overline{B \ker\phi} = \ker\phi$. If one of the followings hold*

(i) *$B$ is not left $\phi$-amenable;*

(ii) *$X$ has a right $\phi$-character;*

*then $T^{**}$ is not approximately biflat.*

*Proof.* We go toward a contradiction and suppose that $T^{**}$ is approximately biflat. Let $\psi_\phi$ be same as above. It is clear that $\ker\psi_\phi = Tri(A, X, \ker\phi)$. Since $\overline{A^2} = A$, $\overline{A \cdot X} = X$ and $\overline{B \ker\phi} = \ker\phi$, then $\overline{T \ker\psi_\phi} = \psi_\phi$. Then by Theorem 3.5, $T$ is left $\psi_\phi$-amenable. Set $I = Tri(0, X, B)$. It is clear that $I$ is closed ideal and $\psi_\phi|_I \neq 0$. Using [8, Lemma 3.1], one can see that $I$ is left $\psi_\phi$-amenable. Thus by [8, Theorem 1.4], there exists a bounded net $(i_\alpha)$ in $I$ such that

$$ii_\alpha - \psi_\phi(i)i_\alpha \to 0, \quad \psi_\phi(i_\alpha) = 1, \quad (i \in I).$$

Take $(x_\alpha)$ in $X$ and $(b_\alpha)$ in $B$ such that $i_\alpha = \begin{pmatrix} 0 & x_\alpha \\ 0 & b_\alpha \end{pmatrix}$. Hence we have

$$\begin{pmatrix} 0 & x \\ 0 & b \end{pmatrix} \begin{pmatrix} 0 & x_\alpha \\ 0 & b_\alpha \end{pmatrix} - \psi_\phi(\begin{pmatrix} 0 & x \\ 0 & b \end{pmatrix}) \begin{pmatrix} 0 & x_\alpha \\ 0 & b_\alpha \end{pmatrix} \to 0$$

and $\psi_\phi(\begin{pmatrix} 0 & x_\alpha \\ 0 & b_\alpha \end{pmatrix}) = \phi(b_\alpha) = 1$ for each $x \in X, b \in B$. Thus we have

$$xb_\alpha - \phi(b)x_\alpha \to 0, bb_\alpha - \phi(b)b_\alpha \to 0, \phi(b_\alpha) = 1, \quad (x \in X, b \in B).$$

If $(i)$ holds the facts

$$bb_\alpha - \phi(b)b_\alpha \to 0, \phi(b_\alpha) = 1, \quad (x \in X, b \in B)$$

give us a contradiction(left $\phi$-amenability condition for $B$).

Suppose that $(ii)$ happens. Take $\eta$ as a right $\phi$-character on $X$. Since $xb_\alpha - \phi(b)x_\alpha \to 0, \phi(b_\alpha) = 1$ for each $x \in X, b \in B$, then we have

$$\eta(xb_\alpha - \phi(b)x_\alpha) = \eta(xb_\alpha) - \phi(b)\eta(x_\alpha) = \phi(b_\alpha)\eta(x) - \phi(b)\eta(x_\alpha) \to 0$$

for each $x \in X, b \in B$. Thus we have $\lim \phi(b)\eta(x_\alpha) = \eta(x)$. Take $b \in \ker\phi$, then we have $\eta(x) = 0$ for each $x \in X$ which is a contradiction($\eta$ is a non-zero functional). □

**Corollary 5.2.** *Let $G$ be a locally compact group. Then $Tri(S(G), L^1(G), S(G))^{**}$ is not approximately biflat.*

*Proof.* Suppose that $\phi \in \Delta(S(G))$. It is well-known that $S(G)$ has a left approximate identity and is a dense left ideal of $L^1(G)$ and also $L^1(G)$ has a bounded approximate identity. Then

$$\overline{S(G)^2} = S(G), \quad \overline{S(G)L^1(G)} = L^1(G), \quad \overline{S(G)\ker\phi} = \ker\phi.$$

Since $S(G)$ is a left ideal in $L^1(G)$, by [2, Lemma 2.2] $\phi$ can be extended to a character on $L^1(G)$, which is a right $\phi$-character for $L^1(G)$. Now apply Theorem 5.1 to show that $Tri(S(G), L^1(G), S(G))^{**}$ is not approximately biflat. □

**Corollary 5.3.** *Let $G$ be a locally compact group. Then $Tri(L^1(G), S(G) \otimes_p S(G), S(G))^{**}$ is not approximately biflat.*

*Proof.* Since $L^1(G)$ has a bounded approximate identity, we have $\overline{L^1(G)^2} = L^1(G)$. Also using Cohn factorization theorem we have $\overline{L^1(G) \cdot (S(G) \otimes_p S(G))} = S(G) \otimes_p S(G)$. Since $S(G)$ has a left approximate identity, then $\overline{S(G)\ker\phi} = \ker\phi$ for each $\phi \in \Delta(S(G))$. Note that for each $\phi \in \Delta(S(G))$, $\phi \otimes \phi$ which is defined by

$$\phi \otimes \phi(a \otimes b) = \phi(a)\phi(b) \quad (a, b \in S(G))$$

is a right $\phi$-character on $S(G) \otimes_p S(G)$. Then apply Theorem 5.1 to show that $Tri(L^1(G), S(G) \otimes_p S(G), S(G))^{**}$ is not approximately biflat. □

Similarly one can show the following result.

**Corollary 5.4.** *Let $G$ be a locally compact group. Then $Tri(L^1(G), M(G), S(G))^{**}$ is not approximately biflat.*

## References


[1] H. P. Aghababa, L. Y. Shi and Y. J. Wu, *Generalized notions of character amenability* Act. Math. Sin, **29** (2013) 1329-1350.

[2] M. Alaghmandan, R. Nasr Isfahani and M. Nemati; *Character amenability and contractibility of abstract Segal algebras*, Bull. Aust. Math. Soc, **82** (2010) 274-281.

[3] M. Essmaili, M. Rostami, A. Pourabbas; *Pseudo-amenability of certain semigroup algebras*, Semigroup Forum (2011), 478-484.

[4] F. Ghahramani,R. J. Loy, G. A. Willis; *Amenability and weak amenability of second conjugate Banach algebras*, Proc. Amer. Math. Soc. **124** (1996), 1489-1497.

[5] F. Ghahramani, Y. Zhang; *Pseudo-amenable and pseudo-contractible Banach algebras*, Math. Proc. Camb. Phil. Soc. **142** (2007) 111-123.

[6] A. Ya. Helemskii; *The homology of Banach and topological algebras*, Kluwer, Academic Press, Dordrecht, 1989.

[7] J. Howie; *Fundamental of Semigroup Theory*. London Math. Soc Monographs, vol. **12**. Clarendon Press, Oxford (1995).

[8] E. Kaniuth, A. T. Lau and J. Pym; *On $\phi$-amenablity of Banach algebras*, Math. Proc. Cambridge Philos. Soc. **144** (2008) 85-96.

[9] J. L. Kelley; *General topology*, D. Van Nostrand Company, Inc., Toronto-New York- London, 1955.

[10] A. R. Medghalchi, M. H. Sattari; *Biflatness and biprojectivity of triangular Banach algebras*. Bull. Iran. Math. Soc. **34**, 115-120 (2008).

[11] P. Ramsden;*Biflatness of semigroup algebras*. Semigroup Forum **79**, (2009) 515-530.

[12] H. Reiter; *$L^1$-algebras and Segal Algebras*, Lecture Notes in Mathematics **231** (Springer, 1971).

[13] E. Samei, N. Spronk, R. Stokke; *Biflatness and pseudo-amenability of Segal algebras*, Canad. J. Math. **62** (2010), 845-869.

Faculty of Mathematics and Computer Science, Amirkabir University of Technology, 424 Hafez Avenue, 15914 Tehran, Iran.

*E-mail address*: amir.sahami@aut.ac.ir